

Mean Convergence of Vector-valued Walsh Series

Jörg Wenzel

Abstract

Given any Banach space X , let L_2^X denote the Banach space of all measurable functions $f : [0, 1] \rightarrow X$ for which

$$\|f\|_2 := \left(\int_0^1 \|f(t)\|^2 dt \right)^{1/2}$$

is finite. We show that X is a UMD-space (see [1]) if and only if

$$\lim_n \|f - S_n(f)\|_2 = 0 \quad \text{for all } f \in L_2^X,$$

where

$$S_n(f) := \sum_{i=0}^{n-1} \langle f, w_i \rangle w_i$$

is the n -th partial sum associated with the Walsh system (w_i) .

1 Introduction

There are several enumerations of the system of Walsh functions. Therefore we first give the appropriate definition.

For $i \geq 1$, the Rademacher functions (r_i) are defined as follows

$$r_1(t) := \begin{cases} +1 & \text{for } t \in [0, \frac{1}{2}) + \mathbb{Z} \\ -1 & \text{for } t \in [\frac{1}{2}, 1) + \mathbb{Z} \end{cases} \quad \text{and} \quad r_{i+1}(t) := r_1(2^i t).$$

Let $n \in \mathbb{N}$. Then n has a unique representation of the form

$$n = \sum_{i=0}^{\infty} n_i 2^i,$$

with $n_i \in \{0, 1\}$. Note that in fact only a finite number of the n_i are different from zero. We let

$$w_n(t) := \prod_{i=0}^{\infty} r_{i+1}(t)^{n_i}.$$

Again the formally infinite product on the righthand side is finite, hence w_n is well defined.

For $f \in L_2^X$, we denote by $S_n(f)$ the n -th partial sum of the Walsh series of f ,

$$S_n(f) := \sum_{i=0}^{n-1} \langle f, w_i \rangle w_i,$$

where

$$\langle f, w_i \rangle := \int_0^1 f(t) w_i(t) dt.$$

Let X and Y be Banach spaces. For $T : X \rightarrow Y$, the ideal norm $\delta(T|\mathcal{W}_n, \mathcal{W}_n)$ is defined to be the least constant $c \geq 0$ such that for all $f \in L_2^X$

$$\|TS_n(f)\|_2 \leq c\|f\|_2.$$

Note that

$$\delta(T|\mathcal{W}_{2^p}, \mathcal{W}_{2^p}) = \|T\| \tag{1}$$

for all operators T and $p \in \mathbb{N}$; see e. g. [2]. In order to get a non-decreasing sequence of ideal norms, we let

$$\delta^{\max}(T|\mathcal{W}_n, \mathcal{W}_n) := \max_{1 \leq k \leq n} \delta(T|\mathcal{W}_k, \mathcal{W}_k).$$

For a more general treatment of ideal norms associated with orthogonal systems we refer to [5], from where the above notation is adopted.

For $k = 1, \dots, 2^n$, let

$$\Delta_k^{(n)} := \left[\frac{k-1}{2^n}, \frac{k}{2^n} \right)$$

be the k -th dyadic intervall of order n .

A dyadic martingale is a martingale $(M_0, M_1, \dots)$ relative to the dyadic filtration $\mathcal{F} = (\mathcal{F}_n)$, where $\mathcal{F}_n$ is generated by $\{\Delta_k^{(n)} : k = 1, \dots, 2^n\}$. If $(M_0, M_1, \dots)$ is an X -valued dyadic martingale, then there exist elements $x_j \in X$ such that

$$M_i = \sum_{j=0}^{2^i-1} x_j h_j, \tag{2}$$

where h_j denotes the j -th Haar function

$$h_0 \equiv 1$$

$$h_j(t) := \begin{cases} +2^{(p-1)/2} & \text{for } t \in \Delta_{2^{m+1}}^{(p)}, \\ -2^{(p-1)/2} & \text{for } t \in \Delta_{2^{m+2}}^{(p)}, \\ 0 & \text{otherwise,} \end{cases}$$

and $j = 2^{p-1} + m$, $m = 0, \dots, 2^{p-1} - 1$.

As usual, we let

$$dM_i := M_{i+1} - M_i.$$

Given $p \in \{1, 2, \dots\}$, let $\mu_p(T)$ denote the least constant $c \geq 0$ such that for all X -valued dyadic martingales $(M_0, M_1, \dots, M_p)$ and for all sequences $\varepsilon_0, \dots, \varepsilon_{p-1}$ of signs ± 1 we have

$$\left\| \sum_{i=0}^{p-1} \varepsilon_i T dM_i \right\|_2 \leq c \|M_p\|_2.$$

We write $\mu_p(X)$ instead of $\mu_p(I_X)$, where I_X denotes the identity map of the Banach space X .

Note that for all $T : X \rightarrow Y$

$$\mu_{p-1}(T) \leq \mu_p(T). \quad (3)$$

Choosing $M_p := M_{p-1}$ in the defining inequality of $\mu_p(T)$, we get $dM_{p-1} = 0$ and hence

$$\left\| \sum_{i=0}^{p-2} \varepsilon_i T dM_i \right\|_2 \leq \mu_p(T) \|M_{p-1}\|_2,$$

which proves the desired inequality.

With the above notation we can prove the following result.

Theorem *For all operators $T : X \rightarrow Y$ and $p \in \mathbb{N}$, we have*

$$\delta^{\max}(T|\mathcal{W}_{2^p}, \mathcal{W}_{2^p}) \leq \mu_p(T) \leq 2\delta^{\max}(T|\mathcal{W}_{2^p}, \mathcal{W}_{2^p}).$$

By definition a Banach space X has the UMD-property if there exists a constant $c \geq 0$ such that

$$\left\| \sum_{i=0}^n \varepsilon_i dM_i \right\|_2 \leq c \left\| \sum_{i=0}^n dM_i \right\|_2$$

for all martingales $(M_0, M_1, \dots)$ with values in X and all $n \in \mathbb{N}$. This is equivalent to the boundedness of the sequence $\mu_p(X)$; see [1].

Thus the theorem gives a characterization of UMD-spaces by the mean convergence of X -valued Walsh series.

2 Preliminaries

Let $s, t \in [0, 1]$. Then s and t have unique representations $s = \sum_{j=0}^{\infty} s_j 2^{-j-1}$ and $t = \sum_{j=0}^{\infty} t_j 2^{-j-1}$, respectively, supposed we choose them to be finite if possible. By $s \oplus t$ we denote the dyadic sum of s and t ,

$$s \oplus t := \sum_{j=0}^{\infty} |t_j - s_j| 2^{-j-1}.$$

Then

$$\int_0^1 f(s) ds = \int_0^1 f(s \oplus t) ds \quad (4)$$

for all $f \in L_1$ and $t \in [0, 1]$. Moreover

$$w_n(s \oplus t) = w_n(s)w_n(t). \quad (5)$$

For $n \geq 1$, let

$$D_n(t) := \sum_{i=0}^{n-1} w_i(t)$$

be the n -th Dirichlet kernel associated with the Walsh functions.

We have

$$S_n(f)(t) = \int_0^1 f(s) D_n(s \oplus t) ds. \quad (6)$$

For $n \geq 1$, let $0 \leq k_1 < k_2 < \dots < k_s$ be defined by

$$n = \sum_{l=1}^s 2^{k_l}. \quad (7)$$

We will use the following result from [6, Theorem 8, p. 28].

Lemma 1

$$D_n = w_n \sum_{i \in \{k_1, \dots, k_s\}} (D_{2^{i+1}} - D_{2^i}).$$

3 Proof of the theorem

For n as in (7), we have by (5), (6) and lemma 1

$$\begin{aligned} \|TS_n(f)\|_2 &= \left\| \sum_{i \in \{k_1, \dots, k_s\}} (TS_{2^{i+1}}(fw_n) - TS_{2^i}(fw_n)) \right\|_2 \\ &\leq \frac{1}{2} \left\| \sum_{i=0}^{k_s} (TS_{2^{i+1}}(fw_n) - TS_{2^i}(fw_n)) \right\|_2 + \\ &\quad + \frac{1}{2} \left\| \sum_{i=0}^{k_s} \varepsilon_i (TS_{2^{i+1}}(fw_n) - TS_{2^i}(fw_n)) \right\|_2, \end{aligned} \quad (8)$$

where ε_i is defined by

$$\varepsilon_i := \begin{cases} +1 & \text{if } i \in \{k_1, \dots, k_s\} \\ -1 & \text{if } i \notin \{k_1, \dots, k_s\} \end{cases}.$$

Note that $M_i := S_{2^i}(fw_n)$ form a dyadic martingale of the form (2), since the linear span of the Walsh functions $w_0, \dots, w_{2^p-1}$ coincides with the linear span of the Haar functions $h_0, \dots, h_{2^p-1}$; see [4]. Hence we have

$$\begin{aligned} \left\| \sum_{i=0}^{k_s} \varepsilon_i (TS_{2^{i+1}}(fw_n) - TS_{2^i}(fw_n)) \right\|_2 &= \left\| \sum_{i=0}^{k_s} \varepsilon_i TdM_i \right\|_2 \\ &\leq \mu_{k_s+1}(T) \|S_{2^{k_s+1}}(fw_n)\|_2. \end{aligned}$$

The same argument applied with $\varepsilon_i = +1$ for all $i = 0, \dots, k_s$ yields

$$\begin{aligned} \left\| \sum_{i=0}^{k_s} (TS_{2^{i+1}}(fw_n) - TS_{2^i}(fw_n)) \right\|_2 &= \left\| \sum_{i=0}^{k_s} TdM_i \right\|_2 \\ &\leq \mu_{k_s+1}(T) \|S_{2^{k_s+1}}(fw_n)\|_2. \end{aligned}$$

Therefore we obtain from (8) that

$$\|TS_n(f)\|_2 \leq \mu_{k_s+1}(T) \|S_{2^{k_s+1}}(fw_n)\|_2.$$

If $n < 2^p$, then $k_s + 1 \leq p$ and it follows from (1) and (3) that

$$\|TS_n(f)\|_2 \leq \mu_{k_s+1}(T) \|fw_n\|_2 \leq \mu_p(T) \|f\|_2.$$

If $n = 2^p$, then again by (1)

$$\|TS_n(f)\|_2 \leq \|T\| \|S_n(f)\|_2 \leq \mu_p(T) \|f\|_2.$$

Consequently

$$\|TS_n(f)\|_2 \leq \mu_p(T) \|f\|_2$$

for all $1 \leq n \leq 2^p$ and hence

$$\delta^{\max}(T|\mathcal{W}_{2^p}, \mathcal{W}_{2^p}) \leq \mu_p(T).$$

This proves the lefthand inequality of the theorem.

To check the righthand inequality we use the following fact.

Lemma 2 *Let $I \subseteq \{0, \dots, p-1\}$ and let n be defined by $n := \sum_{i \in I} 2^i < 2^p$. Then we have*

$$\left\| \sum_{i \in I} TdM_i \right\|_2 \leq \delta(T|\mathcal{W}_n, \mathcal{W}_n) \|M_p\|_2$$

for all martingales $(M_0, \dots, M_p)$ of the form (2).

PROOF: We write M_i in the form

$$M_i = \sum_{j=0}^{2^i-1} x_j w_j,$$

where

$$x_j := \int_0^1 M_p(t) w_j(t) dt \in X.$$

Then, by lemma 1,

$$\begin{aligned} \left\| \sum_{i \in I} T dM_i \right\|_2 &= \left\| \sum_{i \in I} \left(\sum_{j=2^i}^{2^{i+1}-1} T x_j w_j \right) \right\|_2 \\ &= \left\| \sum_{i \in I} (T S_{2^{i+1}}(M_p) - T S_{2^i}(M_p)) \right\|_2 \\ &= \left\| \sum_{i \in I} \left(\int_0^1 T M_p(s) (D_{2^{i+1}}(s \oplus t) - D_{2^i}(s \oplus t)) ds \right) \right\|_2 \\ &= \left\| \int_0^1 T M_p(s) w_n(s \oplus t) D_n(s \oplus t) ds \right\|_2 \\ &= \|T S_n(M_p w_n)\|_2 \\ &\leq \delta(T|\mathcal{W}_n, \mathcal{W}_n) \|M_p w_n\|_2 \\ &= \delta(T|\mathcal{W}_n, \mathcal{W}_n) \|M_p\|_2. \end{aligned}$$

□

We are now able to complete the proof of the theorem. Let a sequence $\varepsilon_0, \dots, \varepsilon_{p-1}$ of signs ± 1 be given. Define n and m by

$$n := \sum_{\{i: \varepsilon_i = +1\}} 2^i \quad \text{and} \quad m := \sum_{\{i: \varepsilon_i = -1\}} 2^i.$$

Then we get from lemma 2 that

$$\begin{aligned} \left\| \sum_{i=0}^{p-1} \varepsilon_i T dM_i \right\|_2 &\leq \left\| \sum_{\{i: \varepsilon_i = +1\}} T dM_i \right\|_2 + \left\| \sum_{\{i: \varepsilon_i = -1\}} T dM_i \right\|_2 \\ &\leq \delta(T|\mathcal{W}_n, \mathcal{W}_n) \|M_p\|_2 + \delta(T|\mathcal{W}_m, \mathcal{W}_m) \|M_p\|_2 \\ &\leq 2\delta^{\max}(T|\mathcal{W}_{2^p}, \mathcal{W}_{2^p}) \|M_p\|_2. \end{aligned}$$

Since this holds for all sequences (ε_i) , we have

$$\mu_p(T) \leq 2\delta^{\max}(T|\mathcal{W}_{2^p}, \mathcal{W}_{2^p}),$$

which is the desired righthand inequality.

4 Some consequences

Corollary 1 *The following conditions are equivalent.*

- (i) $\|f - S_n(f)\|_2 \rightarrow 0$ for all $f \in L_2^X$.
- (ii) X has the UMD-property.

PROOF: If X has the UMD-property, then by the theorem

$$\delta^{\max}(X|\mathcal{W}_n, \mathcal{W}_n) \leq c$$

for all $n \in \mathbb{N}$. Since the Walsh functions form a complete orthonormal system in $L_2[0, 1]$, we can find a linear combination $\sum_{k=0}^N x_k w_k \in L_2^X$ with

$$\left\| f - \sum_{k=0}^N x_k w_k \right\|_2 \leq \varepsilon.$$

Then, for $n \geq N$,

$$\begin{aligned} \|f - S_n(f)\|_2 &\leq \left\| f - \sum_{k=0}^N x_k w_k \right\|_2 + \left\| S_n\left(\sum_{k=0}^N x_k w_k - f\right) \right\|_2 \\ &\leq \varepsilon + \delta^{\max}(X|\mathcal{W}_n, \mathcal{W}_n)\varepsilon \leq (1 + c)\varepsilon. \end{aligned}$$

If on the other hand

$$\|f - S_n(f)\|_2 \rightarrow 0 \quad \text{for all } f \in L_2^X,$$

then, by the uniform boundedness theorem, we get

$$\|S_n(f)\|_2 \leq c\|f\|_2$$

and hence X is a UMD-space. $\square$

As a further easy application of the theorem we get the order of growth of $\mu_p(X)$ for $X = L_1[0, 1]$.

To this end, let

$$L_n := \int_0^1 |D_n(t)| dt \tag{9}$$

be the Lebesgue constants associated with the Walsh system. We also consider

$$L_n^{\max} := \max_{k \leq n} L_k.$$

Corollary 2

$$\frac{p}{8} \leq \mu_p(L_1[0, 1]) \leq 2p.$$

PROOF: For $X = L_1[0, 1]$ and $f \in L_1[0, 1]$, we define $F \in L_2^X$ by

$$F(t) := f(t \oplus \cdot).$$

Then, by (4) and (5), we have

$$\begin{aligned} \langle F, w_j \rangle &= \int_0^1 F(t) w_j(t) dt = \int_0^1 f(t \oplus \cdot) w_j(t) dt \\ &= \int_0^1 f(t) w_j(t) w_j(\cdot) dt = \langle f, w_j \rangle w_j. \end{aligned}$$

Hence

$$\sum_{j=0}^{n-1} \langle F, w_j \rangle w_j(t) = \sum_{j=0}^{n-1} \langle f, w_j \rangle w_j(t \oplus \cdot)$$

Furthermore

$$\begin{aligned} \|F\|_2^2 &= \int_0^1 \|F(t)\|_1^2 dt = \int_0^1 \left(\int_0^1 |f(t \oplus s)| ds \right)^2 dt \\ &= \int_0^1 \left(\int_0^1 |f(s)| ds \right)^2 dt = \|f\|_1^2. \end{aligned}$$

Similarly

$$\left\| \sum_{j=0}^{n-1} \langle F, w_j \rangle w_j \right\|_2 = \left\| \sum_{j=0}^{n-1} \langle f, w_j \rangle w_j \right\|_1.$$

If we now choose f to be the characteristic function of the interval $[0, 2^{-p}]$, then for $k \leq n < 2^p$

$$\langle f, w_k \rangle = 2^{-p} \quad \text{and} \quad \|f\|_1 = 2^{-p}.$$

Consequently

$$\begin{aligned} \delta(L_1 | \mathcal{W}_n, \mathcal{W}_n) &\geq \frac{\left\| \sum_{j=0}^{n-1} \langle F, w_j \rangle w_j \right\|_2}{\|F\|_2} \\ &= L_n, \end{aligned}$$

where L_n denotes the Lebesgue constant as defined in (9).

Since

$$\delta(X | \mathcal{W}_n, \mathcal{W}_n) \leq L_n$$

for all Banach spaces X , we have

$$\delta(L_1 | \mathcal{W}_n, \mathcal{W}_n) = L_n.$$

This proves corollary 2 by taking into account our theorem and the following result from [6, Theorem 9, p. 34].

Lemma 3

$$\frac{p}{8} \leq L_{2^p}^{\max} \leq p.$$

□

References

- [1] Burkholder, D. L.: Martingales and Fourier analysis in Banach spaces. In *Probability and analysis (Varenna, Italy, 1985)*, Lecture Notes in Mathematics, 1206, pages 61–108. Springer–Verlag, 1986.
- [2] Голубов, Б. И.; Ефимов, А. В.; Скворцов, Ш. А.: *Ряды и преобразования Уолша*. Наука, Москва, 1987.
- [3] Кашин, Б. С.; Саакян, А. А.: *Ортогональные Ряды*. Наука, Москва, 1984.
- [4] Paley, R. E. A. C.: A remarkable series of orthogonal functions I. *Proceedings of the London Mathematical Society*, 34:241–264, 1932.
- [5] Pietsch, A.; Wenzel, J.: Orthogonal systems and geometry of Banach spaces. in preparation.
- [6] Schipp, F.; Wade, W. R.; Simon, P.: *Walsh series: an introduction to dyadic harmonic analysis*. Adam Hilger, Bristol, New York, 1990.

Mathematische Fakultät
Friedrich–Schiller–Universität Jena
Universitätshochhaus 17. OG
O–6902 Jena
Germany
e-mail: wenzel@mathematik.uni-jena.de.dbp (X400)